

# Quantum Fluctuations and Irreversibility<sup>1</sup>.

J. Pérez–Mercader<sup>2</sup>

Laboratorio de Astrofísica Espacial y Física Fundamental  
Apartado 50727  
28080 Madrid

February 1, 1993

## ABSTRACT

We study how the effects of quantum corrections lead to notions of irreversibility and clustering in quantum field theory. In particular, we consider the virtual “charge” distribution generated by quantum corrections and adopt for it a statistical interpretation. Then, this virtual charge is shown to (*a*) describe a system where the equilibrium state is at its classical limit ( $\hbar \rightarrow 0$ ), (*b*) give rise to spatial diffusion of the virtual cloud that decays as the classical limit is approached and (*c*) lead to a scenario where clustering takes place due to quantum dynamics, and a natural transition from a “fractal” to a homogeneous regime occurs as distances increase.

---

<sup>1</sup>To be published in “The Physical Origins of Time Asymmetry”, Eds., J. Halliwell, J. Pérez–Mercader and W. Zurek, Cambridge University Press (1993).

<sup>2</sup>Also at: Instituto de Matemáticas y Física Fundamental, C.S.I.C., Serrano 119–123, 28006 Madrid and the Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545.

In quantum field theory, the dynamics stems from both the interactions among the fields and the quantum fluctuations to which they are subject. Quantum fluctuations are perhaps the most fundamental feature of quantum field theory, affecting fields, their sources and the vacuum in which they evolve.

Induced by the fact that quantum fluctuations are to a certain extent random <sup>[1]</sup>, we will study how this affects irreversibility. Namely, since *true* randomness impairs one’s ability to carry out an exact reconstruction of the past history of the system, it is legitimate to expect some irreversibility at the microscopic (quantum field) level because of the presence of quantum fluctuations in this domain.

One possible way to study this irreversibility is to introduce some fundamental statistical notions into the realm of the quantum field theory. For example, one can consider the effect that quantum fluctuations have on the “charge” density<sup>3</sup> of the Poisson equation satisfied by the quantum corrected potentials. The quantum fluctuations affect the interaction energy, and this induces a modified “charge” density; this is how the Uehling potential appears in QED. The statistical notions come in when we interpret the induced charge density as a probability density, an action that we are justified in taking due to the mathematical properties of the charge density induced by quantum–fluctuations. We associate with the full interaction energy a “charge” density which (as is done in cosmology and astrophysics<sup>[2]</sup>) we interpret as a probability density characterizing the spatial distribution of virtual charges surrounding the original charges. By doing this, we are making use of the rich *conceptual* depth of quantum field theory and explicitly taking into account that, in the computation of the quantum corrections at some scale, we are actually including not only simple two body processes with a fixed and precise impact parameter of the order of that scale, but the effects of many body interactions with impact parameter smaller than the distance scale that we are probing. Because of this, divergences appear in the theory which renormalization removes and re–interprets. The physical potential that one measures is thus subject to fluctuations, and it is impossible to specify the *exact*

---

<sup>3</sup>We enclose the word charge in quotes because we refer to it in a generic sense, that is, we are not specifically referring to electric charge. We have in mind the charge for the source of the Poisson equation satisfied by the effective interaction energy.

dependence of the potential on the individual components of the virtual “charge” density. We may, however, ask about the probability of occurrence of some particular configuration or similar questions of a statistical nature. This goal is achieved by adopting for the virtual “charge” charge density a statistical interpretation in term of a probability distribution whose density is the quantum corrected “charge” density. By studying the *effective* (in the RG <sup>[3]</sup> sense) form of the interaction energy, one acquires information on how the properties of the virtual cloud change with scale and, through the statistical interpretation, gain insight on how changes of scale affect the dynamics of the cloud in aspects relating to the approach to equilibrium or even structure and form generation properties.

We can obtain the effective interaction energy by solving its RGE. The solution to this equation<sup>[4]</sup> yields the interaction energy as a function of the scale at which one probes the system, and contains the modifications due to the presence of quantum fluctuations. Unfortunately, the corrections that are included to *all* orders of perturbation theory are only the leading logarithms; but, as is well known, even with this limitation, many interesting physical consequences can still be extracted. Since energy has canonical dimension of inverse time and no anomalous dimension, the RGE has the solution,

$$V(\lambda r_0, g_0, a) = \lambda^{-1} V(r_0, \bar{g}_0(\lambda), a) \quad (1)$$

Here  $V$  is the interaction energy,  $r_0$  is the distance between the interacting sources, and  $a$  is a reference distance. The quantity  $\bar{g}_0(\lambda)$  is the effective coupling and  $\lambda$  the scale parameter;  $\bar{g}_0(\lambda)$  satisfies the RGE  $\lambda \partial \bar{g}_0 / \partial \lambda = -\beta(\bar{g}_0)$ .

In general, for a massless mediating field (such as photons or gluons), the effective interaction energy for two point particles separated by a distance  $a$  is given by

$$V(a, g_0, a) = C \frac{g_0^2}{4\pi a} \quad (2)$$

Computing to 1-loop order, where  $\beta = \beta_0 g_0^3$ , choosing  $\lambda = r/a$ , and taking  $\sigma$  to be a constant<sup>4</sup> (for  $r$  close to  $a$ )

---

<sup>4</sup>Strictly speaking,  $\sigma$  is a function of the distance at which the system is probed. This can be

$$V(r, g_0, a) = C \frac{g_0^2}{4\pi} a^{-\sigma} r^{-1+\sigma} \quad (3)$$

$\sigma$  is related to the  $\beta$ -function for the coupling  $g_0$ , and to one-loop is given by  $\sigma = -2\beta_0 g_0^2$ . It has three essential properties: (i) it is proportional to  $\hbar$ , (ii) it vanishes smoothly in the limit of zero coupling and (iii) as defined, it is a function of the momenta which essentially counts the number of degrees of freedom excited from the vacuum at distances less than  $r_0$ .

For the case of QED one recognizes here the first term in the short distance expansion of the famous Uehling potential; in QCD one recognizes the expression for the interquark potential.

According to potential theory<sup>[6]</sup>, Equation (3) satisfies a Poisson equation whose right hand side is proportional to the “charge” density dictated by quantum corrections. This charge density describes how the phenomenon of vacuum polarization has modified the vacuum and given rise to the formation of virtual pairs that, as is well known, affect the strength and properties of the interaction in a substantial way.

However, we have no information as to the actual distribution of these pairs, but by interpreting the vacuum polarization charge density as a “probability density”, we can derive information on the virtual cloud and its physics as a many body (statistical) system. From the mathematical point of view this is possible thanks to the intimate relationship<sup>[6]</sup> that exists between potential theory and probability theory. The relationship between the potential and the charge density (away from  $r = 0$ ) is through the Poisson equation  $\nabla^2 \phi = +4\pi \rho(r)$ , which for our isotropic potential gives

$$\rho(r) = A' r^{-3+\sigma} \quad (4)$$

where  $A'$  is a constant. We only need to require from the density  $\rho(r)$  that it be a positive and integrable function on its support in order that it can actually be interpreted as a probability density (see below).

---

made explicit by computing it in a mass dependent subtraction procedure<sup>[5]</sup>. However, it will be sufficient for our present purposes to take it to be a constant.

We now examine some properties of Eq.(4). First of all we see that  $\rho(r)$  is the solution to the functional equation

$$\rho(\lambda r) = \lambda^\beta \rho(r) \quad (5)$$

with  $\beta = -3 + \sigma$ ; this implies right away that  $\rho(r)$  describes a fractal distribution of charge embedded in 3-dimensional configuration space, and with a Hausdorff (or fractal) dimension given by  $d_f = +\sigma$ . This is not surprising, since  $\sigma$  has its origin in the deviation from *canonical* scaling due to the quantum fluctuations. Furthermore, since  $\sigma$  changes as we change the size of the domain on which we probe the virtual cloud, it turns out then that the Hausdorff dimension also changes and we are then dealing with a multifractal. For QCD and Quantum Gravity in its asymptotically free regime,  $\sigma$  is positive. In QED and other non-asymptotically free theories,  $\sigma$  is negative. When the Hausdorff dimension is positive, one sees intuitively that there is a natural tendency to suppress these fluctuations since, contrary to what happens in the case of a negative Hausdorff dimension, the configuration space “cannot accommodate them”: it is not big enough!

To reveal some of the physical consequences of the randomness associated with these quantum fluctuations, we will study a few of the features of the “statistical mechanics” of the density in Eq.(4). In particular, we will look at the entropy associated with  $\rho$  and also will give a glance at the nature of the stochastic processes that it supports.

Requiring normalizability of the probability density, we get

$$\rho = A r^{-3+\sigma} \quad (6)$$

with

$$A = \frac{\sigma}{4\pi} R_0^{-\sigma} \quad (7)$$

for  $\sigma > 0$ , and

$$A = -\frac{\sigma}{4\pi} r_0^{-\sigma} \quad (8)$$

when  $\sigma < 0$ . Here  $R_0$  denotes an IR cutoff, necessary in the case of positive  $\sigma$ , in order to ensure the finiteness of the probability distribution. For non-asymptotically free theories,  $r_0$  is an UV cutoff which becomes necessary for the same reasons. The IR cutoff can be identified, e. g., with a typical hadronic size and  $r_0$  with Planck's length. The resulting probability density is of the Pareto type<sup>[7]</sup>; this is a natural consequence of the renormalization group origin for  $\rho(r)$ , which is ultimately responsible for the functional equation in (5) and the associated scaling. Scaling leads to Levy-type distributions, which turn Pareto in some limit. Notice also that both densities go to zero in the classical ( $\hbar \rightarrow 0$ ) limit.

To gain information about the “equilibrium configurations” supported by these distributions, we construct and compute an entropy-like quantity. We will assume that, as in thermodynamics, the state of equilibrium corresponds to the maximum for the entropy. In other words, we will assume that it reveals a preferred “direction” for stability in the evolution of the quantum field system. It is possible to discuss two types of entropies: a “coarse grained” entropy based on the probability distribution, and a “fine grained” entropy or “differential” entropy<sup>5</sup>, based on the probability density  $\rho(r)$ . We will discuss only the latter, which is defined through

$$S = -k \int d^3\vec{r} \rho(r) \log [C\rho(r)] \quad (9)$$

with the integral extended over the full support of the variable  $r$ . The constant  $C$  needs to be introduced because  $\rho(r)$  is a dimensional quantity, and we do not have the equivalent of a Nernst theorem to set a reference valid for *all* physical systems. This constant, as in thermodynamics,<sup>[9]</sup> fixes the minimum entropy of the system. We will set Boltzmann's constant  $k$  equal to one.

The differential entropy for the case when  $\sigma$  is positive gives

---

<sup>5</sup>In the same sense as in information theory<sup>[8]</sup>.

$$S^{(\sigma>0)} = 1 - \frac{3}{\sigma} - \log \frac{\sigma}{4\pi} C + 3 \log R_0 \quad (10)$$

and for negative  $\sigma$  one gets,

$$S^{(\sigma<0)} = 1 - \frac{3}{\sigma} - \log \frac{-\sigma}{4\pi} C + 3 \log r_0 \quad (11)$$

From these two expressions we see that the “entropy constant”,  $C$ , may be related in an interesting and useful way to the “extreme” volume (the volume at the cutoff) of the quantum system. Such a choice has the advantage of eliminating the dependence of the entropy on the cutoff. For asymptotically free theories, if we choose  $C$  proportional to the maximum volume, i.e., proportional to the volume of the IR-cutoff to the cube, then the dependence on the cutoff and extreme system size disappears from the expression for the entropy. The equivalent statement is also true for the case of non-asymptotically free theories, where the “Nernst-volume” corresponds to the minimum volume that can be physically reachable.

These entropies are shown in Figures 1 and 2.

We see that in asymptotically free theories, the equilibrium state corresponds to the *largest* possible value<sup>6</sup> of  $\sigma$ . Now, since matter fluctuations contribute negatively to  $\sigma$ , the most stable state corresponds to a configuration (or system size) where the matter states do not contribute. Because of the decoupling theorem<sup>[10]</sup>, this occurs for the *maximum-possible size* of the system: that is for its *classical* limit!

On the other hand, when the theory is non-asymptotically free, the maximum of the entropy happens when  $\sigma$  goes to 0 from the left. This means either the classical limit, in the sense that  $\hbar \rightarrow 0$ , or that one probes the system at distances *so large* that no quantum fluctuation contributes to  $\sigma$ .

Thus, for both, asymptotically-free and non-asymptotically-free theories one sees that *the maximum of the entropy is attained at the classical limit*.

How does this transition to the equilibrium state take place? Since diffusion is a mechanism for the transition to the equilibrium state for a system where randomness

---

<sup>6</sup>For positive  $\sigma$ ,  $S^{(\sigma>0)}$  has a maximum at  $\sigma = 3$ ; unfortunately, this value lies outside the range where we can reasonably trust the approximations made here.

is at work, we will examine diffusion in quantum field theory, and check if the emerging picture is consistent with what we have obtained from the entropy. In quantum field theory, the fractal nature of the probability density leads us to expect some form of fractal diffusion or, more precisely, diffusion through fractal brownian motion. Because of this “fractality” we *expect* the diffusion to be non-space filling, and to lead to either clustering ( $\sigma > 0$ ) or screening when the system relaxes; it is clear that this behavior ought to depend on the sign of  $\sigma$ . In what follows we will examine some of the properties of the coefficient of diffusion derived from  $\rho(r)$ ; then we will examine the correlation integral in some specific cases.

The virtual charge density cloud described by the probability density Equation (6) will “diffuse” in a manner similar<sup>[11]</sup> to what occurs in Brownian motion. We expect that the components of the virtual cloud scatter randomly among themselves; the collision probability density is  $\rho(r)$ ; the virtual charges execute random walks controlled by  $\rho(r)$ ; the net effect of these processes is that the virtual cloud diffuses outside of the ball of radius  $R$ .

As is well known<sup>[11]</sup> the probability that a “particle” executing a random walk, with each individual step governed by a probability density  $\rho(r)$ , be after  $N$ -steps at a position between  $\vec{r}$  and  $\vec{r} + d\vec{r}$ , is given by

$$W(r)d^3\vec{r} = [4\pi Dt]^{-3/2} \exp \left[ -|\vec{r}|^2 / (4Dt) \right] d^3\vec{r} \quad (12)$$

where  $D$  is the coefficient of diffusion,

$$D = \frac{n}{6} \langle \vec{r}^2 \rangle ,$$

$n$  is the number of steps per unit time and  $N = nt$  is the total number of steps.  $\langle \vec{r}^2 \rangle$  is the second moment of the probability density  $\rho(r)$ . The function  $W(r)$  in Eq. (12) satisfies the 3-dimensional diffusion equation

$$\frac{\partial W}{\partial t} = D \nabla^2 W$$



with boundary condition that  $W|_s = 0$  at an absorbing medium, or vanishing normal derivative of  $W$  at a reflecting surface:  $\frac{\partial W}{\partial n}|_s = 0$ . How far and how fast a disturbance propagates depends on the coefficient of diffusion: for large  $D$  one has large amounts of diffusion, and viceversa. When  $D = 0$ , there is no diffusion.

In addition to the the Markovian nature of the process, there are two assumptions involved in the derivation of (12) which we must bear in mind: (a) the number of steps  $N$  is very large and (b) in the computation of the characteristic function, it is assumed that  $kr \ll 1$ . These two assumptions mean in our case that the time intervals that we consider are “long” compared with virtual time intervals and that the size of the region in which we expect the diffusion to take place, is “large” compared to the Compton wavelength about which we are computing the coefficient  $\sigma$ . Both of these restrictions are amply met by the validity of our approximations.

In order to compute the coefficient of diffusion one needs to obtain the second moment of the probability density. From a radial distance  $s_0$  to a radial distance  $s > s_0$ , the second moment of  $\rho(r)$  is

$$\langle r^2 \rangle = \frac{4\pi}{2+\sigma} A (s^{2+\sigma} - s_0^{2+\sigma}) \quad (13)$$

and therefore, in some cases, the coefficient of diffusion *diverges* (naively) as  $s$  goes to infinity. This again is not surprising, since our probability distribution is the limit of a Levy-type distribution, and thus its moments are divergent.

As previously, we classify the situation depending on whether we have an asymptotically free theory or not. When  $\sigma > 0$ , we can take  $s_0 = 0$  and set  $s = R_0$ , the IR cutoff. The coefficient of diffusion is given by

$$D^{(\sigma>0)} = \frac{n}{6} \frac{\sigma}{2+\sigma} R_0^2. \quad (14)$$

In non-asymptotically free theories, we take  $s_0 = r_0$  (the UV-cutoff) and leave  $s$  unspecified. Then,

$$D^{(\sigma<0)} = \frac{n}{6} \frac{-\sigma}{2+\sigma} s^2 \left[ \left( \frac{s}{r_0} \right)^\sigma - \left( \frac{r_0}{s} \right)^2 \right]. \quad (15)$$

The expression for  $D^{(\sigma>0)}$  reflects the divergence in the second moment of Levy-type distributions. However, quantum corrections again play an important rôle here: physics limits the size of the system ( $R_0$  cannot go to infinity) and in the classical limit, when  $\sigma \rightarrow 0$ , diffusion *stops*. Before this happens, i.e., when the typical energies are larger than the inverse Compton wavelength of the IR-cutoff, the system will tend to diffuse itself into clusters or well differentiated pieces. This last statement being a consequence of the positive fractal (Hausdorff) dimension associated with  $\sigma > 0$ , as was mentioned above.

For  $\sigma < 0$  the situation is qualitatively different. We can distinguish two different regimes within non-asymptotically free theories, according as to whether  $0 > \sigma > -2$  or  $-2 > \sigma$ . In the former case,  $\langle \vec{r}^2 \rangle$  diverges as diffusion takes place into larger distances, and is only shut-off by the vanishing of  $\sigma$  with distance, that is to say, by the transition into the classical regime. When  $\sigma < -2$ , the coefficient of diffusion goes to zero as  $s$  increases; this means that *new* states stationary under diffusion can be created and supported at these very deep values of  $\sigma$ . When  $\sigma > -2$  the diffusion coefficient diverges for large  $s$  and quantum corrections will shut the diffusion process off by eventually driving  $\sigma$  to zero; what happens is that as  $s$  increases, the diffusion coefficient at first increases, to then decrease and cut itself off to zero when  $\sigma$  goes to zero or when the classical regime ( $\hbar \rightarrow 0$ ) is reached.

We have seen how quantum corrections lead into diffusion, and in some cases, clustered states of the quantum field system. To get more information about clustering and other structure formation properties, it is convenient to look at the correlation function and integral; from an analysis of these objects one can get a better feeling of the structures<sup>[12]</sup> supported by the quantum field theory. The correlation integral,  $C(s)$ , gives information on how many correlated pairs there are whose separation is *less* than  $s$ . This integral is computed from the correlation function  $\xi(\vec{r})$  by the following formula<sup>[13],[2]</sup>

$$C(s) = \int_{V(s)} d^3\vec{r} [1 + \xi(\vec{r})]$$

where the region of integration extends over the volume  $V(s)$  of radius  $s$ . The correlation function  $\xi(\vec{r})$  is related to  $\delta(\vec{k})$ , the Fourier transform of the density  $\rho(\vec{r})$ , via the following Fourier transform

$$1 + \xi(\vec{r}) = (2\pi)^3 \int d^3\vec{k} e^{-i\vec{k}\cdot\vec{r}} |\delta(\vec{k})|^2$$

where

$$\delta(\vec{k}) = \int d^3\vec{r} e^{+i\vec{k}\cdot\vec{r}} \rho(\vec{r})$$

The quantity  $|\delta(\vec{k})|^2$  is called the power spectrum, and measures the mean number of correlated neighbors in *excess* of random over a distance of order  $1/k$ ; because of this, the quantity  $1 + \xi(\vec{r})$  (once appropriately normalized) gives information on whether there is “clustering” ( $> 1$ ), “voids” ( $< 1$ ) or a perfectly random distribution ( $= 1$ ). This is a straightforward program that can be carried out for  $\rho(r)$ . One obtains the following expressions for  $1 + \xi(\vec{r})$  and  $C(r)$

$$1 + \xi(r) = D(\sigma) \cdot r^{-3+\sigma}$$

$$C(r) = \frac{2\pi}{\sigma} \cdot D(\sigma) \cdot r^{2\sigma}$$

where the coefficient  $D(\sigma)$  is

$$D(\sigma) = (4\pi A)^2 \cdot 4\pi \cdot |B(\sigma)|^2 \cdot \Gamma(2 - 2\sigma) \cos \frac{\pi}{2}(2\sigma - 1)$$

and

$$B(\sigma) = \Gamma(\sigma - 1) \cos \frac{\pi}{2}(2 - \sigma).$$

$D(\sigma)$  is shown in Figure 3, and in Figure 4 we show  $D(\sigma)/\sigma$ , which are important in determining the behaviors of  $1 + \xi(r)$  and the correlation integral.

From Figure 3 we see that for positive  $\sigma$  and between zero and  $3/2$ , the quantum field system leads to clustering. For  $\sigma$  between  $3/2$  and  $5/2$  there is a significant qualitative change into a scenario of deep anticlustering (“voids”) with a tendency for the system to behave near homogeneity when  $\sigma \approx 2$ . When  $\sigma$  increases past  $5/2$ , we enter a new clustering domain. The rest can be “read off” in the same way from the figure. The plot of Figure 4 simply confirms the one for  $D(\sigma)$ , but for the coefficient of the correlation integral. We also point out that there is a “quantization” of structures, a kind of periodicity, that takes place as  $\sigma$  changes and goes through some critical values.

We end by summarizing our results. Quantum field theory is a many body system par excellence. Intrinsic to it is the randomness associated with quantum fluctuations. The application of a statistical interpretation to one of the most primitive concepts of field theory, that of the induced charge, has interesting and apparently deep consequences which stem from its scaling properties. In particular, fractal behavior in a form related to the Pareto–Levy form of the charge density, leads to systems where the maximum entropy is associated with the classical limit of the quantum system. In other words, the classical regime is the most stable, and the one preferred by the system.

Like in other many body systems, diffusion is a familiar mechanism for the approach to equilibrium (although perhaps not the only one). In concordance to what follows from studying the entropy, the virtual cloud diffuses with a coefficient of diffusion proportional to  $\hbar$ , and thus diffusion of the cloud stops in the classical limit.

From an analysis of the correlation properties of the virtual cloud system, one learns that the system supports the formation of structures where there is a sharp transition between phases of anticlustering (void formation), clustering and some interspersed quasi–homogeneity. These depend quite clearly on the number of degrees of freedom excited from the vacuum into the system; in other words, they depend on the distance at which the system is probed.

In very general terms, we see that irreversibility is contained in quantum field systems, and that this irreversibility is encountered as the system size grows. The

quantum field system tends to “relax” into a classical system or into clustered structures, depending on the scale of the system and the nature of the interactions. The ideas developed here may also find application in the study of intermittency, turbulence, phase transitions, multiparticle physics and other complex phenomena in quantum field theory and the early universe.

## Acknowledgements.

This workshop was financed by Fundación Banco de Bilbao Vizcaya and by NATO. The Fundación also provided us with logistic, material and public support. I wish to thank Don José Angel Sánchez Asiaín, Doña Maria Luisa Oyarzábal and Don José Ignacio Oyarzábal for their enthusiastic support. I would also like to thank my co-organizers, Jonathan Halliwell and Wojciech Zurek for the pleasure and fun in working together. I have benefitted from discussions with O. Bertolami, R. Blankebeckler, A. Carro, T. Goldman, A. González–Arroyo, S. Habib, J. Hartle, R. Laflamme, S. Lloyd, C. Morales, M. M. Nieto, E. Trillas, G. Veneziano and G. West.

## References

- [1] Bell, J. S., ., *Speakeable and unspeakeable in quantum mechanics*, Cambridge University Press, Cambridge, 1991.
- [2] See for example, Layzer, D., in *Galaxies and the Universe*, edited by Sandage, A., Sandage, M. and Kristian, J., University of Chicago Press, Chicago, 1975.
- [3] Gell–Mann, M. and Low, F. E., *Phys. Rev.* 95 (1954) 1300–1312. For a beautiful presentation full of insight, see West, G. in *Particle Physics: A Los Alamos Primer*, edited by Cooper, N. and West, G., Cambridge University Press, Cambridge, 1988.
- [4] Rothe, H. J., *Lattice Gauge Theories*, World Scientific, Singapore, 1992.
- [5] Georgi, H. and Politzer, D., *Phys. Rev.* D14 (1974) 451.
- [6] Doob, J., ., *Classical Potential Theory and Its Probabilistic Counterpart*, Springer–Verlag, New York, 1984.
- [7] Schroeder, M., *Fractals, Chaos, Power Laws*, W. H. Freeman and Co., San Francisco, 1991.
- [8] Cover, T. M. and Thomas, J. A., *Elements of Information Theory*, J. Wiley and Sons, New York, 1991.
- [9] Fermi, E., *Thermodynamics*, Dover Books, New York, 1956.

- [10] Appelquist, T. and Carrazone, J., *Phys. Rev. D*11 (1975) 2896.
- [11] Chandrasekhar, S., *Rev. Mod. Phys.*15 (1943) 1–89.
- [12] Thompson, D’Arcy Wentworth, *On Growth and Form: The Complete Revised Edition*, Dover Books, New York, 1992.
- [13] Grassberger, P. and Procaccia, I. *Physica D*9 (1983) 189.